

Graphs with large generalized 3-edge-connectivity*

Xueliang Li¹, Yaping Mao¹, Lusheng Wang²

¹Center for Combinatorics and LPMC-TJKLC, Nankai University
Tianjin 300071, China. Email: lxl@nankai.edu.cn; maoyaping@ymail.com

²Department of Computer Science, City University of Hong Kong
Tat Chee Avenue, Kowloon, Hong Kong SAR, China. Email: lwang@cs.cityu.edu.hk

Abstract

Let G be a graph, and S a set of vertices of G . Denote by $\lambda(S)$ the maximum number ℓ of pairwise edge-disjoint trees $T_1, T_2, \dots, T_\ell$ in G such that $V(T_i) \supseteq S$ for every $1 \leq i \leq \ell$. Then the generalized k -edge-connectivity $\lambda_k(G)$ of G is defined as $\lambda_k(G) = \min\{\lambda(S) | S \subseteq V(G) \text{ and } |S| = k\}$. Thus $\lambda_2(G) = \lambda(G)$. In this paper, sharp upper and lower bounds of $\lambda_3(G)$ are given for a connected graph G of order n , that is, $1 \leq \lambda_3(G) \leq n - 2$. Graphs of order n such that $\lambda_3(G) = n - 2, n - 3$ are characterized, respectively.

Keywords: edge-connectivity, edge-disjoint trees, generalized edge-connectivity.

AMS subject classification 2010: 05C40, 05C05.

1 Introduction

All graphs considered in this paper are undirected, finite and simple. We refer to the book [1] for graph theoretical notation and terminology not described here.

The generalized connectivity of a graph G , which was introduced by Chartrand et al. in [2], is a natural and nice generalization of the concept of (vertex) connectivity. Let G be a nontrivial connected graph of order n and $2 \leq k \leq n$. For a set S of k vertices of G , let $\kappa(S)$ denote the maximum number ℓ of edge-disjoint trees $T_1, T_2, \dots, T_\ell$ in G such that $V(T_i) \cap V(T_j) = S$ for every pair i, j of distinct integers with $1 \leq i, j \leq \ell$ (Note that the trees are vertex-disjoint in $G \setminus S$). A collection $\{T_1, T_2, \dots, T_\ell\}$ of trees in G with this property is called an *internally disjoint set of trees connecting S* . The *generalized k -connectivity*, denoted by $\kappa_k(G)$, of G is then defined as $\kappa_k(G) = \min\{\kappa(S) | S \subseteq V(G) \text{ and } |S| = k\}$. Thus, $\kappa_2(G) = \kappa(G)$.

As a natural counterpart of the generalized connectivity, recently we introduced the concept of generalized edge-connectivity in [9]. Let $\lambda(S)$ denote the maximum number ℓ of pairwise edge-disjoint trees $T_1, T_2, \dots, T_\ell$ in G such that $V(T_i) \supseteq S$ for every $1 \leq i \leq \ell$. Then the *generalized k -edge-connectivity* $\lambda_k(G)$ of G is defined as $\lambda_k(G) = \min\{\lambda(S) | S \subseteq V(G) \text{ and } |S| = k\}$. Thus $\lambda_2(G) = \lambda(G)$.

*Supported by NSFC No.11071130, and RGCHK Project No. CityU 9041382.

In addition to being natural combinatorial measures, the generalized connectivity and generalized edge-connectivity can be motivated by their interesting interpretation in practice. For example, suppose that G represents a network. If one considers to connect a pair of vertices of G , then a path is used to connect them. However, if one wants to connect a set S of vertices of G with $|S| \geq 3$, then a tree has to be used to connect them. This kind of tree with minimum order for connecting a set of vertices is usually called a Steiner tree, and popularly used in the physical design of VLSI, see [11]. Usually, one wants to consider how tough a network can be, for connecting a set of vertices. Then, the number of totally independent ways to connect them is a measure for this purpose. The generalized k -connectivity and generalized k -edge-connectivity can serve for measuring the capability of a network G to connect any k vertices in G .

There have been many results on the generalized connectivity, see [2, 3, 10, 5, 6, 7, 8]. In [9] we obtained some results on the generalized edge-connectivity. The following results are restated, which will be used later.

Lemma 1. [9] *For every two integers n and k with $2 \leq k \leq n$, $\lambda_k(K_n) = n - \lceil k/2 \rceil$.*

Lemma 2. [9] *For any connected graph G , $\lambda_3(G) \leq \lambda(G)$. Moreover, the upper bound is sharp.*

In [4], we characterized graphs with large generalized 3-connectivity and obtained the following result.

Lemma 3. [4] *Let G be a connected graph of order n . Then $\kappa_3(G) = n - 2$ if and only if $G = K_n$ or $G = K_n \setminus e$.*

Like [4], here we will consider the generalized 3-edge-connectivity. In Section 3, sharp upper and lower bounds of $\lambda_3(G)$ are given for a connected graph G of order n , that is, $1 \leq \lambda_3(G) \leq n - 2$. Graphs of order n such that $\lambda_3(G) = n - 2$, $n - 3$ are characterized, respectively.

2 Preliminary

For a graph G , let $V(G)$ and $E(G)$ denote the set of vertices and the set of edges of G , respectively, and $|G|$ and $\|G\|$ the order and the size of G , respectively. If M is a subset of edges of a graph G , the subgraph of G induced by M is denoted by $G[M]$. As usual, the *union* of two graphs G and H is the graph, denoted by $G \cup H$, with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. Let mH be the disjoint union of m copies of a graph H . For $W \subseteq V(G)$, we denote $G \setminus W$ the subgraph by deleting the vertices of W together with the edges incident with them from G . If S is a subset of vertices of a graph G , the subgraph of G induced by S is denoted by $G[S]$. If M is the edge subset of G , then $G \setminus M$ denote the subgraph by deleting the edges of M . If $M = e$, we simply write $G \setminus e$ for $G \setminus \{e\}$. We denote by $E[X, Y]$ the set of edges of G with one end in X and the other end in Y . If $X = \{x\}$, we simply write $E[x, Y]$ for $E[\{x\}, Y]$. A subset M of $E(G)$ is called a *matching* of G if the edges of M satisfy that no two of them are adjacent in G . A matching M saturates a vertex v , or v is said to be *M -saturated*, if some edge of M is

incident with v ; otherwise, v is M -unsaturated. M is a *maximum matching* if G has no matching M' with $|M'| > |M|$.

Lemma 4. *Let G be a connected graph with minimum degree δ . Then $\lambda_3(G) \leq \delta$. In particular, if there are two adjacent vertices of degree δ , then $\lambda_3(G) \leq \delta - 1$.*

Proof. We know that $\lambda(G) \leq \delta$ and $\lambda_3(G) \leq \lambda(G)$ by Lemma 2. So $\lambda_3(G) \leq \delta$.

Suppose that there are two adjacent vertices v_1 and v_2 of degree δ and $\lambda_3(G) = \delta$. Besides v_1 and v_2 , we choose a vertex v_3 in $V(G \setminus \{v_1, v_2\})$ to get a set $S = \{v_1, v_2, v_3\}$. Suppose $T_1, T_2, \dots, T_\delta$ are δ pairwise edge-disjoint trees connecting S . Since G is simple graph, obviously the δ edges incident v_1 must be contained in $T_1, T_2, \dots, T_\delta$, respectively, and so are the δ edges incident v_2 . Without loss of generality, we may assume that the edge v_1v_2 is contained in T_1 . But, since T_1 is a tree connecting S , it must contain another edge incident with v_1 or v_2 , a contradiction. Thus $\lambda_3(G) \leq \delta - 1$. $\square$

Lemma 5. *If G is a graph obtained from the complete graph K_n by deleting an edge set M and $\Delta(K_n[M]) \geq 3$, then $\lambda_3(G) \leq n - 4$.*

Proof. Suppose that $\lambda_3(G) \geq n - 3$. Since $\Delta(K_n[M]) \geq 3$, there exists at least one vertex, denoted by v , such that $d_{K_n[M]}(v) \geq 3$. Then $d_G(v) = n - 1 - d_{K_n[M]}(v) \leq n - 4$. So $\delta(G) \leq d_G(v) \leq n - 4$. From Lemma 4, $\lambda_3(G) \leq \delta(G) \leq n - 4$, a contradiction. $\square$

Lemma 5 indicates that if $\lambda_3(G) \geq n - 3$, then each component of $K_n[M]$ must be a path or a cycle (Note that an independent edge is also a path of length 1).

Now we state a useful Lemma for general k , although we only consider the case for $k = 3$ in this paper.

Let $S = \{u_1, u_2, \dots, u_k\} \subseteq V(G)$ and $W = V(G) \setminus S = \{w_1, w_2, \dots, w_{n-k}\}$. Then let A be a maximum set of edge-disjoint trees in G that connect S . Let A_1 be the set of trees in A whose edges belong to $E(G[S])$, and let A_2 be the set of trees containing at least one edge of $E[S, W]$. Thus $A = A_1 \cup A_2$. For a tree T connecting S in G , if $T \in A_1$, we call it a tree of Type I ; if $T \in A_2$, we call it a tree of Type II .

Lemma 6. *Let $S \subseteq V(G)$, $|S| = k$ and T be a tree connecting S . If $T \in A_1$, then T uses $k - 1$ edges of $E(G[S]) \cup E[S, W]$; If $T \in A_2$, then T uses at least k edges of $E(G[S]) \cup E[S, W]$.*

Proof. It is easy to see that for each T in A_1 , T uses $k - 1$ edges in $E(G[S])$, namely, T uses $k - 1$ edges of $E(G[S]) \cup E[S, W]$.

For $T \in A_2$, by deleting all the vertices of T in W , we obtain some components of T in S , denoted by $C_1, C_2, \dots, C_s$. Let $|C_i| = k_i$. Then $\|C_i\| = k_i - 1$ and $\sum_{i=1}^s (k_i - 1) = k - s$. Since there exists at least one edge of T between each C_i and W , where $1 \leq i \leq s$, T uses at least $(k - s) + s = k$ edges in $E(G[S]) \cup E[S, W]$. $\square$

3 Graphs with 3-edge-connectivity $n - 2, n - 3$

The following observation is easily seen from definition.

Observation 1. If G be a connected graph, then $\kappa_k(G) \leq \lambda_k(G) \leq \delta(G)$.

After the preparation of the above section, we start to give our main results of this paper.

Theorem 1. For a connected graph G of order n , $1 \leq \lambda_3(G) \leq n - 2$. Moreover, the upper and lower bounds are sharp.

Proof. It is easy to see that $\lambda_3(G) \leq \lambda_3(K_n)$. From this together with Lemma 1, we have $\lambda_3(G) \leq n - 2$. Since G is connected, then $\lambda_3(G) \geq 1$. The result holds.

One can easily check that the complete graph K_n attains the upper bound and the complete bipartite graph $K_{1,n-1}$ attains the lower bound. $\square$

The following result is very similar to Lemma 3 for the generalized 3-connectivity, but the proof for the edge case is different.

Theorem 2. For a connected graph G of order n , $\lambda_3(G) = n - 2$ if and only if $G = K_n$ or $G = K_n \setminus e$, where $e \in E(G)$.

Proof. If $G = K_n$, then $\lambda_3(G) = n - 2$ by Lemma 1. If $G = K_n \setminus e$, then it follows by Observation 1 and Lemma 3 that $\lambda_3(G) \geq \kappa_3(G) = n - 2$. From this together with Theorem 1, $\lambda_3(G) = n - 2$.

On the other hand, we will show that if $G \neq K_n, K_n \setminus e$, then $\lambda_3(G) \leq n - 3$. Actually, we only need to show that $\lambda_3(G) \leq n - 3$ for a graph G obtained from the complete graph K_n by deleting any two edges. Let $G = K_n \setminus \{e_1, e_2\}$, where $e_1, e_2 \in E(K_n)$. It is easy to see that e_1 and e_2 form a path of order 3 (Figure 1 (a)), or e_1 and e_2 are independent edges (Figure 1 (b)). First, we consider the former case. Let $P_3 = x, y, z$, $S = \{x, y, z\}$, and $W = G \setminus S$. Then $|E(G[S]) \cup E[S, W]| = 3(n - 3) + 1$. Since $xy, yz \notin E(G)$, there exists no tree of Type I. So each tree connecting S must belong to Type II. From Lemma 6, each tree of Type II uses at least 3 edges in $E(G[S]) \cup E[S, W]$. So $3(n - 3) + 1$ edges form at most $\frac{3(n-3)+1}{3}$ trees. Thus $\lambda_3(G) \leq |A| = |A_1| + |A_2| = 0 + |A_2| \leq \frac{3(n-3)+1}{3}$ and $\lambda_3(G) = n - 3$ since $\lambda_3(G)$ is an integer.




Figure 1. The two cases for $G = K_n \setminus \{e_1, e_2\}$.

Next we consider the latter case. Let $e_1 = xy$, $e_2 = zw_1$, $S = \{x, y, z\}$, and $W = G \setminus S$. So $w_1 \in W$ and $|E(G[S]) \cup E[S, W]| = 3(n - 3) + 1$. If xz and yz form a tree in A_1 , then the tree use 2 edges in $E(G[S]) \cup E[S, W]$ and the remaining $3(n - 3) - 1$ edges in $E(G[S]) \cup E[S, W]$ form at most $\frac{3(n-3)-1}{3}$ trees of Type II. So $|A| = |A_1| + |A_2| \leq 1 + \frac{3(n-3)-1}{3}$ and $\lambda_3(G) \leq |A| \leq n - 3$ since $\lambda_3(G)$ is an integer. If xz and yz do

not form a tree in A_1 , then all the edges of $E(G[S]) \cup E[S, W]$ can only form a tree of Type II connecting S . From Lemma 6, each tree of Type II uses at least 3 edges in $E(G[S]) \cup E[S, R]$. Thus $\lambda_3(G) \leq |A| = |A_2| \leq \frac{3(n-3)+1}{3}$ and $\lambda_3(G) \leq n-3$. $\square$

For the generalized 3-connectivity, we got the following result in [4].

Theorem 3. [4] *Let G be a connected graph of order n ($n \geq 3$). Then $\kappa_3(G) = n-3$ if and only if G is a graph obtained from the complete graph K_n by deleting an edge set M such that $K_n[M] = P_4$ or $K_n[M] = P_3 \cup P_2$ or $K_n[M] = C_3 \cup P_2$ or $K_n[M] = rP_2$ ($2 \leq r \leq \lfloor \frac{n}{2} \rfloor$).*

But, for the edge case the statement is different.

Theorem 4. *Let G be a connected graph of order n . Then $\lambda_3(G) = n-3$ if and only if G is a graph obtained from the complete graph K_n by deleting an edge set M such that $K_n[M] = rP_2$ ($2 \leq r \leq \lfloor \frac{n}{2} \rfloor$) or $K_n[M] = P_4 \cup sP_2$ ($0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$) or $K_n[M] = P_3 \cup tP_2$ ($0 \leq s \leq \lfloor \frac{n-3}{2} \rfloor$) or $K_n[M] = C_3 \cup tP_2$ ($0 \leq s \leq \lfloor \frac{n-3}{2} \rfloor$).*

Proof. Sufficiency: Assume that $\lambda_3(G) = n-3$. Then $|M| \geq 2$ by Theorem 2 and each component of $K_n[M]$ is a path or a cycle by Lemma 5. We will show that the following claims hold.

Claim 1. $K_n[M]$ has at most one component of order larger than 2.

Suppose, to the contrary, that $K_n[M]$ has two components of order larger than 2, denoted by H_1 and H_2 (See Figure 2). Let $x, y \in V(H_1)$ and $z \in V(H_2)$ such that $d_{H_1}(y) = d_{H_2}(z) = 2$ and x is adjacent to y in H_1 . Pick $S = \{x, y, z\}$. Since $G = K_n \setminus M$, we have $d_G(y) = d_{K_n}(y) - d_{H_1}(y) = n-3$. The same is true for z , that is, $d_G(z) = n-3$. This implies that $\delta(G) \leq n-3$. Since all other components of $K_n[M]$ are paths or cycles, $\delta(G) \geq n-3$. So $\delta(G) = n-3$ and $d_G(y) = d_G(z) = \delta(G)$. Since $yz \in E(G)$, by Lemma 4 it follows that $\lambda_3(G) \leq \delta(G) - 1 = n-4$, a contradiction.




Figure 2. Graphs for Claim 1 (The dotted lines stand for edges in M).

Claim 2. If H is the component of $K_n[M]$ of order larger than 3, then H is a 4-path.

Assume, to the contrary, that H is a path or a cycle of order larger than 4, or a cycle of order 4.

First, we consider the former. We can pick a P_5 in H . Let $P_5 = v_1, v_2, v_3, v_4, v_5$, $S = \{v_2, v_3, v_4\}$ and $W = G \setminus \{v_2, v_3, v_4\}$ (Figure 3 (a)). Since $v_2v_3, v_3v_4 \notin E[S]$, there exists no tree of Type I connecting S . From Lemma 6, each tree of Type II uses at least 3 edges. Since $|E(G[S]) \cup E[S, W]| = 3(n-3) - 1$, we have $|A_2| \leq \frac{3(n-3)-1}{3}$ and $|A| = |A_2| = n-4$ since $\lambda_3(G)$ is an integer. This contradicts to $\lambda_3(G) = n-3$.

Now we consider the latter. Let $H = v_1, v_2, v_3, v_4$ be a cycle, and $S = \{v_2, v_3, v_4\}$ (Figure 3 (b)). Since $v_2v_3, v_3v_4 \notin E[S]$, there exists no tree of Type I. Since each tree of




Figure 3. Graphs for Claim 2.

Type *II* uses at least 3 edges by Lemma 6 and $|E(G[S]) \cup E[S, W]| = 3(n-3) - 1$, we have $|A_2| \leq \frac{3(n-3)-1}{3}$ and $|A| = |A_2| = n-4$, which also contradicts to $\lambda_3(G) = n-3$.

From the above two claims, we know that if $K_n[M]$ has a component P_4 , then it is the only component of order larger than 3 and the other components must be independent edges. Let s be the number of such independent edges. $K_n[M]$ can have as many as such independent edges, which implies that $s \leq \lfloor \frac{n-4}{2} \rfloor$. From Theorem 2, $s \geq 0$. Thus $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$.

By the similar analysis, we conclude that G is a graph obtained from the complete graph K_n by deleting an edge set M such that $K_n[M] = rP_2$ ($2 \leq r \leq \lfloor \frac{n}{2} \rfloor$) or $K_n[M] = P_4 \cup sP_2$ ($0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$) or $K_n[M] = P_3 \cup tP_2$ ($0 \leq t \leq \lfloor \frac{n-3}{2} \rfloor$) or $K_n[M] = C_3 \cup tP_2$ ($0 \leq t \leq \lfloor \frac{n-3}{2} \rfloor$).

Necessity: We will show that $\lambda_3(G) \geq n-3$ if G is a graph with the conditions of this theorem. We have the following cases to consider.

Case 1. $K_n[M] = C_3 \cup tP_2$ ($0 \leq t \leq \lfloor \frac{n-3}{2} \rfloor$) or $K_n[M] = P_3 \cup tP_2$ ($0 \leq t \leq \lfloor \frac{n-3}{2} \rfloor$).

We only need to show that $\lambda_3(G) \geq n-3$ for $t = \lfloor \frac{n-3}{2} \rfloor$. If $\lambda_3(G) \geq n-3$ for $K_n[M] = C_3 \cup \lfloor \frac{n-3}{2} \rfloor P_2$, then $\lambda_3(G) \geq n-3$ for $K_n[M] = P_3 \cup \lfloor \frac{n-3}{2} \rfloor P_2$. So we only need to consider the former. Let $C_3 = v_1, v_2, v_3$ and $S = \{x, y, z\}$ be a 3-set of G and $M_1 = M \setminus E(C_3)$. Since $K_n[M] \setminus V(C_3)$ has $\lfloor \frac{n-3}{2} \rfloor$ independent edges, M_1 is a maximum matching of $G \setminus V(C_3)$. Then $G \setminus V(C_3)$ has at most one M_1 -unsaturated vertex.




Figure 4. Graphs for Case 1.

If $S = V(C_3)$, then there exist $n-3$ pairwise edge-disjoint trees connecting S since each vertex in S is adjacent to every vertex in $G \setminus S$. Suppose $S \neq V(C_3)$.

If $|S \cap V(C_3)| = 2$, then one element of S belongs to $V(G) \setminus V(C_3)$, denoted by z . Since $d_G(v_1) = d_G(v_2) = d_G(v_3) = n-3$, we can assume that $x = v_1, y = v_2$. When z is M_1 -unsaturated, the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$ form $n-3$ pairwise edge-disjoint trees connecting S , where $\{w_1, w_2, \dots, w_{n-4}\} = V(G) \setminus \{x, y, z, v_3\}$.

When z is M_1 -saturated, we let z' be the adjacent vertex of z under M_1 . Then the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$ and $T_2 = xz' \cup yz' \cup z'v_3 \cup zv_3$ form $n-3$ pairwise edge-disjoint trees connecting S (See Figure 4 (a)), where $\{w_1, w_2, \dots, w_{n-5}\} = V(G) \setminus \{x, y, z, z', v_3\}$.

If $|S \cap V(C_3)| = 1$, then two elements of S belong to $\in V(G) \setminus V(C_3)$, denoted by y and z . Without loss of generality, let $x = v_2$. When y and z are adjacent under M_1 , the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xy \cup yv_1 \cup v_1 z$ and $T_2 = xz \cup zv_3 \cup v_3 y$ form $n-3$ pairwise edge-disjoint trees connecting S (See Figure 4 (b)), where $\{w_1, w_2, \dots, w_{n-5}\} = V(G) \setminus \{x, y, z, v_1, v_3\}$. When y and z are nonadjacent under M_1 , we consider whether y and z are M_1 -saturated. If one of $\{y, z\}$ is M_1 -unsaturated, without loss of generality, we assume that y is M_1 -unsaturated. Since $G \setminus V(C_3)$ has at most one M_1 -unsaturated vertex, z is M_1 -saturated. Let z' be the adjacent vertex of z under M_1 . Then the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xy \cup yz$ and $T_2 = v_1 y \cup v_1 z \cup z'v_1 \cup z'x$ and $T_3 = xz \cup zv_3 \cup v_3 y$ form $n-3$ pairwise edge-disjoint trees connecting S (See Figure 4 (c)), where $\{w_1, w_2, \dots, w_{n-6}\} = V(G) \setminus \{x, y, z, z', v_1, v_3\}$. If both y and z are M_1 -saturated, we let y', z' be the adjacent vertex of y, z under M_1 , respectively. Then the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$, $T_2 = xy \cup yz' \cup z'y' \cup y'z$, $T_3 = yv_3 \cup z'v_3 \cup zv_3 \cup xz'$ and $T_4 = yv_1 \cup y'v_1 \cup zv_1 \cup y'x$ form $n-3$ pairwise edge-disjoint trees connecting S (See Figure 4 (d)), where $\{w_1, w_2, \dots, w_{n-7}\} = V(G) \setminus \{x, y, z, y', z', v_1, v_3\}$.

Otherwise, $S \subseteq G \setminus V(C_3)$. When one of $\{x, y, z\}$ is M_1 -unsaturated, without loss of generality, we assume that x is M_1 -unsaturated. Since $G \setminus V(C_3)$ has at most one M_1 -unsaturated vertex, both y and z are M_1 -saturated. Let y', z' be the adjacent vertex of y, z under M_1 , respectively. We pick a vertex x' of $V(G) \setminus \{x, y, y', z, z', v_1, v_2, v_3\}$. When x, y, z are all M_1 -saturated, we let x', y', z' be the adjacent vertex of x, y, z under M_1 , respectively. Then the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_j = xv_j \cup yv_j \cup zv_j$ ($1 \leq j \leq 3$) and $T_4 = xy \cup yx' \cup x'z$ and $T_5 = xy' \cup zy' \cup zy$ and $T_6 = zx \cup xz' \cup z'y$ form $n-3$ pairwise edge-disjoint trees connecting S (See Figure 4 (e)), where $\{w_1, w_2, \dots, w_{n-9}\} = V(G) \setminus \{x, y, z, x', y', z', v_1, v_2, v_3\}$.

From the above discussion, we get that $\lambda(S) \geq n-3$ for $S \subseteq V(G)$, which implies $\lambda_3(G) \geq n-3$. From this together with Theorem 2, we have $\lambda_3(G) = n-3$.

Case 2. $K_n[M] = rP_2$ ($2 \leq r \leq \lfloor \frac{n}{2} \rfloor$) or $K_n[M] = P_4 \cup sP_2$ ($0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$).

We only need to show that $\lambda_3(G) \geq n-3$ for $r = \lfloor \frac{n}{2} \rfloor$ and $s = \lfloor \frac{n-4}{2} \rfloor$. If $\lambda_3(G) \geq n-3$ for $K_n[M] = P_4 \cup \lfloor \frac{n-4}{2} \rfloor P_2$, then $\lambda_3(G) \geq n-3$ for $K_n[M] = \lfloor \frac{n}{2} \rfloor P_2$. So we only need to consider the former. Let $P_4 = v_1, v_2, v_3, v_4$, $S = \{x, y, z\}$ be a 3-set of G , and $M_1 = M \setminus E(P_4)$. Since $K_n[M] \setminus V(P_4)$ has $\lfloor \frac{n-4}{2} \rfloor$ independent edges, M_1 is a maximum matching of $G \setminus V(P_4)$. It is easy to see that $G \setminus V(P_4)$ has at most one M_1 -unsaturated vertex. We will show that there exist $n-3$ trees connecting S in G .

If $S \subseteq V(P_4)$, then there exist $n-4$ pairwise edge-disjoint trees connecting S since each vertex in S is adjacent to every vertex in $G \setminus V(P_4)$. Since $d_G(v_1) = d_G(v_4) = n-2$ and $d_G(v_2) = d_G(v_3) = n-3$, we only need to consider $S = \{v_1, v_2, v_3\}$ and $S = \{v_1, v_2, v_4\}$. These trees together with $T = yv_4 \cup v_4x \cup v_4z$ for $S = \{v_1, v_2, v_3\}$, or $T = xy \cup yz$ for $S = \{v_1, v_2, v_4\}$ form $n-3$ pairwise edge-disjoint trees connecting S . Suppose $S \cap V(P_4) \neq 3$.

If $|S \cap V(P_4)| = 2$, then one element of S belongs to $\in V(G) \setminus V(P_4)$, denoted by z .

Since $d_G(v_1) = d_G(v_4) = n - 2$ and $d_G(v_2) = d_G(v_3) = n - 3$, we only need to consider $x = v_1, y = v_2$ or $x = v_2, y = v_3$ or $x = v_1, y = v_4$. When z is M_1 -unsaturated, the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$, $T_2 = xv_4 \cup yv_4 \cup zv_4$ for $x = v_1, y = v_2$, or $T_2 = xv_4 \cup v_4 v_1 \cup v_1 y \cup v_4 z$ for $x = v_2, y = v_3$, or $T_2 = xv_3 \cup yv_3 \cup zv_3$ for $x = v_1, y = v_4$ form $n - 3$ pairwise edge-disjoint trees connecting S , where $\{w_1, w_2, \dots, w_{n-5}\} = V(G) \setminus (V(P_4) \cup \{z\})$. When z is M_1 -saturated, we let z' be the adjacent vertex of z under M_1 . For $x = v_2, y = v_3$, the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$, $T_2 = xz' \cup yz' \cup z'v_4 \cup zv_4$ and $T_2 = yv_1 \cup v_1 v_4 \cup zv_1 \cup xv_4$ form $n - 3$ pairwise edge-disjoint trees connecting S (See Figure 5 (a)), where $\{w_1, w_2, \dots, w_{n-6}\} = V(G) \setminus \{x, y, z, z', v_1, v_4\}$. One can check that the same is true for $x = v_1, y = v_2$ and $x = v_1, y = v_4$ (See Figure 5 (b) and (c)).




Figure 5. Graphs for S in Case 2.

If $|S \cap V(P_4)| = 1$, then two elements of S belong to $V(G) \setminus V(P_4)$, denoted by y and z . We only need to consider $x = v_1$ or $x = v_2$. When y and z are adjacent under M_1 , the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xy \cup zv_1 \cup yv_1$, $T_2 = xz \cup zv_3 \cup yv_3$ and $T_3 = xv_4 \cup yv_4 \cup zv_4$ form $n - 3$ pairwise edge-disjoint trees connecting S for $x = v_2$ (See Figure 5 (d)), where $\{w_1, w_2, \dots, w_{n-6}\} = V(G) \setminus \{x, y, z, v_1, v_3, v_4\}$. The same is true for $x = v_1$ (See Figure 5 (e)). When y and z are nonadjacent under M_1 , we consider whether y and z are M_1 -saturated. If one of $\{y, z\}$ is M_1 -unsaturated, without loss of generality, we assume that y is M_1 -unsaturated. Since $G \setminus V(P_4)$ has at most one M_1 -unsaturated vertex, z is M_1 -saturated. Let z' be the adjacent vertex of z under M_1 . For $x = v_2$, the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$, $T_2 = v_4 x \cup v_4 y \cup v_4 z$, $T_3 = v_1 y \cup v_1 z \cup xz$ and $T_4 = z'x \cup v_3 y \cup z'v_3 \cup zv_3$ form $n - 3$ pairwise edge-disjoint trees connecting S (See Figure 4 (f)), where $\{w_1, w_2, \dots, w_{n-7}\} = V(G) \setminus \{x, y, z, z', v_1, v_3, v_4\}$. The same is true for $x = v_1$ (See Figure 5 (g)). If both y and z are M_1 -saturated, we let y', z' be the adjacent vertex of y, z under M_1 , respectively. For $x = v_2$, the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_1 = xz \cup yz$, $T_2 = yv_3 \cup zv_3 \cup xz$, $T_3 = xv_4 \cup yv_4 \cup zv_4$, $T_4 = yv_1 \cup y'v_1 \cup zv_1 \cup xy'$ and $T_5 = xz' \cup z'y \cup z'y' \cup y'z$ form $n - 3$ pairwise edge-disjoint trees connecting S (See Figure 5 (h)), where $\{w_1, w_2, \dots, w_{n-8}\} = V(G) \setminus \{x, y, z, y', z', v_1, v_3, v_4\}$. The same is true for $x = v_1$ (See Figure 5 (i)).

If $S \subseteq G \setminus V(P_4)$, when one of $\{x, y, z\}$ is M_1 -unsaturated, without loss of generality, we let x is M_1 -unsaturated, then both y and z are M_1 -saturated. Let y', z' be the adjacent vertex of y, z under M_1 , respectively. We pick a vertex x' of $V(G) \setminus \{x, y, y', z, z', v_1, v_2, v_3\}$. When x, y, z are all M_1 -saturated, we let x', y', z' be the adjacent vertex of x, y, z under M_1 , respectively. Then the trees $T_i = w_i x \cup w_i y \cup w_i z$ together with $T_j = x v_j \cup y v_j \cup z v_j$ ($1 \leq j \leq 4$) and $T_5 = yx \cup xy' \cup y'z$ and $T_6 = yx' \cup zx' \cup zx$ and $T_7 = zy \cup yz' \cup z'x$ form $n - 3$ pairwise edge-disjoint trees connecting S (See Figure 5 (j)), where $\{w_1, w_2, \dots, w_{n-10}\} = V(G) \setminus \{x, y, z, x', y', z', v_1, v_2, v_3, v_4\}$.

The proof is now complete. $\square$

References

- [1] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.
- [2] G. Chartrand, S.F. Kappor, L. Lesniak, D.R. Lick, *Generalized connectivity in graphs*, Bull. Bombay Math. Colloq. 2(1984), 1-6.
- [3] G. Chartrand, F. Okamoto, P. Zhang, *Rainbow trees in graphs and generalized connectivity*, Networks 55(4)(2010), 360-367.
- [4] H. Li, X. Li, Y. Mao, Y. Sun, *Graphs with large generalized 3-connectivity*, arXiv: 1201.2983 [math.CO] 2012.
- [5] S. Li, W. Li, X. Li, *The generalized connectivity of complete bipartite graphs*, Ars Combin. 104(2012).
- [6] S. Li, X. Li, *Note on the hardness of generalized connectivity*, J. Combin. Optimization, in press.
- [7] S. Li, X. Li, Y. Shi, *The minimal size of a graph with generalized connectivity $\kappa_3(G) = 2$* , Australasian J. Combin. 51(2011), 209-220.
- [8] S. Li, X. Li, W. Zhou, *Sharp bounds for the generalized connectivity $\kappa_3(G)$* , Discrete Math. 310(2010), 2147-2163.
- [9] X. Li, Y. Mao, Y. Sun, *The generalized connectivity and generalized edge-connectivity*, arXiv:1112.0127 [math.CO] 2011.
- [10] F. Okamoto, P. Zhang, *The tree connectivity of regular complete bipartite graphs*, J. Combin. Math. Combin. Comput. 74(2010), 279-293.
- [11] N.A. Sherwani, *Algorithms for VLSI Physical Design Automation*, 3rd Edition, Kluwer Acad. Pub., London, 1999.